

Infrared Maximally Abelian Gauge

Tereza Mendes, Attilio Cucchieri and Antonio Mihara

*Instituto de Física de São Carlos, Universidade de São Paulo, Caixa Postal 369,
13560-970 São Carlos, SP, Brazil*

Abstract. The confinement scenario in Maximally Abelian gauge (MAG) is based on the concepts of Abelian dominance and of dual superconductivity. Recently, several groups pointed out the possible existence in MAG of ghost and gluon condensates with mass dimension 2, which in turn should influence the infrared behavior of ghost and gluon propagators. We present preliminary results for the first lattice numerical study of the ghost propagator and of ghost condensation for pure $SU(2)$ theory in the MAG.

Keywords: Yang-Mills theory; Green's functions; Confinement; Abelian projection

PACS: 11.15.Ha 12.38.Aw

The study of the infra-red (IR) limit of QCD is of central importance for understanding the mechanism of confinement. Despite being non-gauge-invariant, gluon and ghost propagators are powerful tools in the (non-perturbative) investigation of this limit. In recent years, (gauge-dependent) condensates of mass dimension two have received considerable attention. An example of such objects is the ghost condensate [1, 2], related to the breakdown of a global $SL(2, R)$ symmetry. In particular, in MAG the diagonal and off-diagonal components of the ghost propagators are expected to be modified by ghost condensation. In this paper we present preliminary results of lattice studies of the Faddeev–Popov (FP) matrix for pure $SU(2)$ theory in the MAG. We consider the ghost propagator, the ghost condensate and the smallest eigenvalue of the FP matrix.

On the lattice, for the $SU(2)$ case, the MAG is obtained (see e.g. [3]) by minimizing the functional

$$S = -\frac{1}{2dV} \sum_{x,\mu} \text{Tr} \left[\sigma_3 U_\mu(x) \sigma_3 U_\mu^\dagger(x) \right]. \quad (1)$$

At any local minimum one has that the Faddeev–Popov matrix, defined as

$$\begin{aligned} \sum_{by} M^{ab}(x,y) \gamma^b(y) &= \sum_{\mu} \gamma^a(x) [V_\mu(x) + V_\mu(x - e_\mu)] + 2 \{ \gamma^a(x - e_\mu) [1 - 2(U_\mu^0(x))^2] \\ &\quad - 2 \sum_b \gamma^b(x - e_\mu) [\varepsilon_{ab} U_\mu^0(x) U_\mu^3(x) + \sum_{cd} \varepsilon_{ad} \varepsilon_{bc} U_\mu^d(x) U_\mu^c(x)] \}, \end{aligned} \quad (2)$$

is positive-definite. Here the color indices take values 1, 2 and we follow the notation $U_\mu(x) = U_\mu^0(x) \mathbb{1} + i \sigma^a U_\mu^a(x)$ and $V_\mu(x) = (U_\mu^0(x))^2 + (U_\mu^3(x))^2 - (U_\mu^1(x))^2 - (U_\mu^2(x))^2$, where σ^a are the 3 Pauli matrices. Notice that (as in Landau gauge [7]) this matrix is symmetric under the simultaneous exchange of color and space-time indices. Using the relation $U_\mu(x) = \exp[-iag_0 A_\mu(x)]$ one finds (in the formal continuum limit $a \rightarrow 0$) the standard continuum results [4] for the stationary conditions above and for $M^{ab}(x,y)$.




FIGURE 1. Left: plot of $G(p^2)$ as a function of improved p for lattice volumes $V = 16^4, 24^4, 40^4$ and $\beta = 2.2$. Right: plot of the smallest eigenvalue of the FP operator, as a function of the inverse linear size of the system.

We have considered four values of β (2.2, 2.3, 2.4, 2.512) and lattice volumes up to 40^4 . Our results for the gluon propagators are in agreement with the study by Bornyakov et al. [3]: we see a clear suppression of the off-diagonal propagators compared to the diagonal (transverse) one, supporting Abelian dominance. We have fitted our data for the various gluon propagators (at all values of V and $\beta = 2.2$), obtaining the following behaviors. For $D(p^2)$ (transverse) diagonal, our data favor a Stingl-Gribov form

$$D(p^2) = \frac{1 + d p^2}{a + b p^2 + c p^4}, \quad (3)$$

with a mass $m = \sqrt{a/b} \approx 0.72 \text{ GeV}$. Note that the above equation corresponds to a pair of complex conjugate poles z and z^* . We can thus write $z = x + iy$ with $x = b/(2c) \approx 0.32 \text{ GeV}^2$ and $y = \sqrt{a/c - x^2} \approx 0.47 \text{ GeV}^2$. Let us recall that in the case of a Gribov-like propagator these two poles are purely imaginary. For $D(p^2)$ transverse off-diagonal our best fit is of Yukawa type, i.e. $D(p^2) = 1/(a + b p^2)$, with a mass $m = \sqrt{a/b} \approx 0.97 \text{ GeV}$. Finally, the longitudinal off-diagonal gluon propagator is best fitted by $D(p^2) = 1/(a + b p^2 + c p^4)$ (i.e. also of Yukawa type) with a mass $m = \sqrt{a/b} \approx 1.25 \text{ GeV}$. As expected from Abelian dominance, the mass is larger in the off-diagonal case.

In Fig. 1 (left) we show our data for the ghost propagator $G(p^2)$, as a function of an improved momentum p (see Ref. [5]). The data show little volume dependence at small p . (Note that, contrary to Landau gauge, here we can evaluate the ghost propagator at zero momentum.) We see no sign of an enhanced IR propagator. We have fitted our data (at $\beta = 2.2$), obtaining a behavior of the type (3) above with $a = 0.45(1) \text{ GeV}^2$, $b = 1.1(3)$, $c = 0.73(30) \text{ GeV}^{-2}$, $d = 2.1(9) \text{ GeV}^{-2}$. Thus, we see a Stingl-Gribov fit with mass $m \approx 0.6 \text{ GeV}$ and complex poles given by $x \approx 0.75$, $y \approx 0.22$.




FIGURE 2. Left: plot of the quantity $\Phi(p^2) = L^2 / \cos(\pi \tilde{p}_\mu a/L) \langle |\varepsilon_{ab} G^{ab}(p^2)/2| \rangle$ as a function of p for lattice volumes $V = 8^4, 16^4, 24^4, 40^4$ and $\beta = 2.2$. Right: plot of $\Phi(p^2)$ as a function of p and L .

We next consider (see Fig. 1, right) the smallest eigenvalue of the FP matrix. We have looked at λ_{min} for several lattice volumes and values of β as a function of $1/L$. The data are fitted to $a(1/L)^b$ with $b = 1.6(1)$, showing that λ_{min} vanishes more slowly than $(1/L)^2$ (Laplacian). This may explain why we do not see a diverging ghost propagator at zero momentum even at rather large lattice volumes [6].

Following the analysis done in Landau gauge [7], we consider the anti-symmetric off-diagonal ghost propagator $\langle |\varepsilon_{ab} G^{ab}(p^2)/2| \rangle$ rescaled by $L^2 / \cos(\pi \tilde{p}_\mu a/L)$, as a function of the (unimproved) momentum p for all lattice volumes and β values considered. The data show nice scaling for all cases considered. The data at $V = 40^4$ and $\beta = 2.2$ can be fitted by $\Phi(p) = (a + b p/L^2)(p^4 + v^2)$ with $a = 0.0026(7) \text{ GeV}^2$, $b = 32.6(7) \text{ GeV}^{-1}$ and $v^2 = 1.7(1) \text{ GeV}^4$. We thus have a rather large ghost condensate $v \approx 1.3 \text{ GeV}^2$, but we cannot be sure that it survives in the infinite-volume limit, since the overall constant a might be null. We can also fit data at several V 's and β 's for $\Phi(p^2)$ as a function of p and L (see Fig. 2, right). We obtain $\Phi(p) = (a + b p/L^2)(p^4 + v^2)$ with $a = 0.0033(6) \text{ GeV}^2$, $b = 35.8(5) \text{ GeV}^{-1}$ and $v^2 = 1.87(8) \text{ GeV}^4$. We note that the fit parameters change little with the (physical) lattice volume.

We are currently investigating the effects of Gribov copies on our results.

We thank M.I. Polikarpov and M. Schaden for helpful discussions. A. C. and T. M. were supported by FAPESP and CNPq. A. M. was supported by FAPESP.

REFERENCES

1. M. Schaden, arXiv:hep-th/9909011.
2. M. A. L. Capri et al., Phys. Rev. D **72**, 085021 (2005).
3. V. G. Bornyakov et al., Phys. Lett. B **559**, 214 (2003).
4. F. Bruckmann, T. Heinzl, A. Wipf and T. Tok, Nucl. Phys. B **584**, 589 (2000).
5. J. P. Ma, Mod. Phys. Lett. A **15**, 229 (2000).
6. A. Cucchieri, these proceedings.
7. A. Cucchieri, T. Mendes and A. Mihara, Phys. Rev. D **72**, 094505 (2005).